

**A major purpose of the Technical Information Center is to provide the broadest dissemination possible of information contained in DOE's Research and Development Reports to business, industry, the academic community, and federal, state and local governments.**

**Although a small portion of this report is not reproducible, it is being made available to expedite the availability of information on the research discussed herein.**

PORTIONS OF THIS REPORT ARE ILLEGIBLE. It  
has been reproduced from the best available  
copy to permit the broadest possible avail-  
ability.

Los Alamos National Laboratory is operated by the University of California for the United States Department of Energy under contract W-7405-ENG-36.

CONF-8405180--2

LA-UR--84-1981

DE84 014032

TITLE: SOME MASSES FOR POPULATION I AND II CEPHEIDS

AUTHOR(S): Russell B. Kidman and Arthur N. Cox, T-6

SUBMITTED TO: Proceedings of IAU Colloquium No. 82, David Dunlap Observatory,  
University of Toronto, May 28 -- June 1, 1984

#### DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

By acceptance of this article, the publisher recognizes that the U.S. Government retains a nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes.

The Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy.

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

**Los Alamos** Los Alamos National Laboratory  
Los Alamos, New Mexico 87545

## SOME MASSES FOR POPULATION I AND II CEPHEIDS

R. B. Kidman <sup>author</sup>

Theoretical Division, Los Alamos National Laboratory, University of California

A. N. Cox

Theoretical Division, Los Alamos National Laboratory, University of California

The masses of Cepheids can be obtained in several ways. If a Cepheid luminosity is known from membership in a galactic cluster, the mass-luminosity relation obtained from stellar evolution theory gives its mass. This evolution mass depends slightly on the composition, that is, the mass fraction of helium,  $Y$ , and on the mass fraction of all the heavier elements,  $Z$ , but as we shall see later, the composition dependence is small.

A mass of a Cepheid that is based entirely on pulsation theory is called the pulsation mass. Here the needed observations are: the luminosity, the color (giving a surface effective temperature by use of a conversion formula), and the easily observed pulsation period. In this report the observed period is assumed to always be the fundamental mode for the classical Cepheids. The mass,  $M$ , the pulsation constant,  $Q$ , and the radius,  $R$ , are unknowns that are solved for using the three equations

$$L = 4\pi R^2 \sigma T_e^4, \quad Q = P(M/R^3)^{1/2}, \quad \text{and} \quad Q = Q(M, R, L, T_e).$$

These equations are: the definition of the effective temperature, the period-mean density relation, and an expression fitting the pulsation theory values for the pulsation constant  $Q$  obtained from a large number of models covering a large range of stellar parameters.

A third mass can be derived based on both evolution and pulsation theories. The required observations are only the well determined period and a surface effective temperature. The above three equations, plus the evolution mass-luminosity relation, are used to simultaneously solve for the unknowns  $M$ ,  $R$ ,  $L$ , and  $Q$ . Due to the strong influence of the evolution theory mass-luminosity relation for the theoretical mass, this mass agrees to within a few % with the evolution mass for almost all Cepheids. The major value of the theoretical mass is its availability when a luminosity has not been observed, and one cannot get an evolution mass.

These three masses have been calculated for 29 Cepheids recently listed by Fernie and McGonegal (1983). The luminosities of these Cepheids have been uniformly set by assuming a distance modulus of 3.29 for the Hyades cluster. Table 1 gives our evolution, theoretical, and pulsation masses. Fernie and McGonegal have suggested that CS Vel and V810 Cen are not in

the cluster as supposed, and therefore their luminosities are incorrectly stated. Due to the large discrepancy between the evolution and pulsation masses for V Cen, GY Sge, and S Vul, we have rejected them also. We find only a small difference between the masses for TW Nor, and therefore we do not reject this Cepheid even though there is a cluster membership question about it. The ratios of these masses relative to the theoretical masses average to the values at the bottom of the evolution and pulsation mass columns. The usual scatter in the pulsation masses is present, but on the average, the masses seem in accord with evolution theory.

Table 2 gives similar results with only the distance scale to the Hyades changed to 3.45, a value recently suggested by Vandenberg and Bridges (1984). With the increased luminosities, evolution masses increase a bit, while the luminosity-independent theoretical masses remain unchanged. However, pulsation masses are greatly increased so that the average is about 1.22 above the theoretical masses. For both these tables we have converted the dereddened (B-V) colors to effective temperatures by use of the Kraft (1961) formula. As discussed by Cox (1979) and many others such as Pel (1978) these temperatures may need to be cooled by 0.01 to 0.03 in  $\log T_e$ . That would greatly alleviate the mass discrepancy. It appears that a distance scale which has the Hyades distance modulus at 3.45 is too large, but part of the change from the currently used 3.29 may be acceptable from the viewpoint of Cepheid masses.

Other recent data on Cepheid luminosities have been prepared for publication by Schmidt (1984). Table 3 shows at the top section the evolution, theoretical, and pulsation masses for 7 Cepheids using his luminosities and dereddened colors. The pulsation masses are so low that for conventional compositions, stars would not evolve into the instability strip. The possible cooling of the temperature scale is investigated in the second section. Even though we have reduced the log

Table 1

Cepheid	Forale and McGonigal Data (m-M) = 3.29					
	P(d)	T (K)	L/L <sub>o</sub>	M <sub>e</sub>	M <sub>p</sub>	M <sub>o</sub>
W Cen	10.1	7000	1.5	4.0	4.0	4.0
W Sge	10.1	7000	1.5	4.0	4.0	4.0
W Vul	10.1	7000	1.5	4.0	4.0	4.0
W Nor	10.1	7000	1.5	4.0	4.0	4.0
W Sge	10.1	7000	1.5	4.0	4.0	4.0
W Vul	10.1	7000	1.5	4.0	4.0	4.0
W Nor	10.1	7000	1.5	4.0	4.0	4.0
W Sge	10.1	7000	1.5	4.0	4.0	4.0
W Vul	10.1	7000	1.5	4.0	4.0	4.0
W Nor	10.1	7000	1.5	4.0	4.0	4.0
W Sge	10.1	7000	1.5	4.0	4.0	4.0
W Vul	10.1	7000	1.5	4.0	4.0	4.0
W Nor	10.1	7000	1.5	4.0	4.0	4.0
W Sge	10.1	7000	1.5	4.0	4.0	4.0
W Vul	10.1	7000	1.5	4.0	4.0	4.0
W Nor	10.1	7000	1.5	4.0	4.0	4.0
W Sge	10.1	7000	1.5	4.0	4.0	4.0
W Vul	10.1	7000	1.5	4.0	4.0	4.0
W Nor	10.1	7000	1.5	4.0	4.0	4.0
W Sge	10.1	7000	1.5	4.0	4.0	4.0
W Vul	10.1	7000	1.5	4.0	4.0	4.0
W Nor	10.1	7000	1.5	4.0	4.0	4.0
W Sge	10.1	7000	1.5	4.0	4.0	4.0
W Vul	10.1	7000	1.5	4.0	4.0	4.0
W Nor	10.1	7000	1.5	4.0	4.0	4.0
W Sge	10.1	7000	1.5	4.0	4.0	4.0
W Vul	10.1	7000	1.5	4.0	4.0	4.0
W Nor	10.1	7000	1.5	4.0	4.0	4.0
W Sge	10.1	7000	1.5	4.0	4.0	4.0
W Vul	10.1	7000	1.5	4.0	4.0	4.0
W Nor	10.1	7000	1.5	4.0	4.0	4.0
W Sge	10.1	7000	1.5	4.0	4.0	4.0
W Vul	10.1	7000	1.5	4.0	4.0	4.0
W Nor	10.1	7000	1.5	4.0	4.0	4.0
W Sge	10.1	7000	1.5	4.0	4.0	4.0
W Vul	10.1	7000	1.5	4.0	4.0	4.0
W Nor	10.1	7000	1.5	4.0	4.0	4.0
W Sge	10.1	7000	1.5	4.0	4.0	4.0
W Vul	10.1	7000	1.5	4.0	4.0	4.0
W Nor	10.1	7000	1.5	4.0	4.0	4.0
W Sge	10.1	7000	1.5	4.0	4.0	4.0
W Vul	10.1	7000	1.5	4.0	4.0	4.0
W Nor	10.1	7000	1.5	4.0	4.0	4.0
W Sge	10.1	7000	1.5	4.0	4.0	4.0
W Vul	10.1	7000	1.5	4.0	4.0	4.0
W Nor	10.1	7000	1.5	4.0	4.0	4.0
W Sge	10.1	7000	1.5	4.0	4.0	4.0
W Vul	10.1	7000	1.5	4.0	4.0	4.0
W Nor	10.1	7000	1.5	4.0	4.0	4.0
W Sge	10.1	7000	1.5	4.0	4.0	4.0
W Vul	10.1	7000	1.5	4.0	4.0	4.0
W Nor	10.1	7000	1.5	4.0	4.0	4.0
W Sge	10.1	7000	1.5	4.0	4.0	4.0
W Vul	10.1	7000	1.5	4.0	4.0	4.0
W Nor	10.1	7000	1.5	4.0	4.0	4.0
W Sge	10.1	7000	1.5	4.0	4.0	4.0
W Vul	10.1	7000	1.5	4.0	4.0	4.0
W Nor	10.1	7000	1.5	4.0	4.0	4.0
W Sge	10.1	7000	1.5	4.0	4.0	4.0
W Vul	10.1	7000	1.5	4.0	4.0	4.0
W Nor	10.1	7000	1.5	4.0	4.0	4.0
W Sge	10.1	7000	1.5	4.0	4.0	4.0
W Vul	10.1	7000	1.5	4.0	4.0	4.0
W Nor	10.1	7000	1.5	4.0	4.0	4.0
W Sge	10.1	7000	1.5	4.0	4.0	4.0
W Vul	10.1	7000	1.5	4.0	4.0	4.0
W Nor	10.1	7000	1.5	4.0	4.0	4.0
W Sge	10.1	7000	1.5	4.0	4.0	4.0
W Vul	10.1	7000	1.5	4.0	4.0	4.0
W Nor	10.1	7000	1.5	4.0	4.0	4.0
W Sge	10.1	7000	1.5	4.0	4.0	4.0
W Vul	10.1	7000	1.5	4.0	4.0	4.0
W Nor	10.1	7000	1.5	4.0	4.0	4.0
W Sge	10.1	7000	1.5	4.0	4.0	4.0
W Vul	10.1	7000	1.5	4.0	4.0	4.0
W Nor	10.1	7000	1.5	4.0	4.0	4.0
W Sge	10.1	7000	1.5	4.0	4.0	4.0
W Vul	10.1	7000	1.5	4.0	4.0	4.0
W Nor	10.1	7000	1.5	4.0	4.0	4.0
W Sge	10.1	7000	1.5	4.0	4.0	4.0
W Vul	10.1	7000	1.5	4.0	4.0	4.0
W Nor	10.1	7000	1.5	4.0	4.0	4.0
W Sge	10.1	7000	1.5	4.0	4.0	4.0
W Vul	10.1	7000	1.5	4.0	4.0	4.0
W Nor	10.1	7000	1.5	4.0	4.0	4.0
W Sge	10.1	7000	1.5	4.0	4.0	4.0
W Vul	10.1	7000	1.5	4.0	4.0	4.0
W Nor	10.1	7000	1.5	4.0	4.0	4.0
W Sge	10.1	7000	1.5	4.0	4.0	4.0
W Vul	10.1	7000	1.5	4.0	4.0	4.0
W Nor	10.1	7000	1.5	4.0	4.0	4.0
W Sge	10.1	7000	1.5	4.0	4.0	4.0
W Vul	10.1	7000	1.5	4.0	4.0	4.0
W Nor	10.1	7000	1.5	4.0	4.0	4.0
W Sge	10.1	7000	1.5	4.0	4.0	4.0
W Vul	10.1	7000	1.5	4.0	4.0	4.0
W Nor	10.1	7000	1.5	4.0	4.0	4.0
W Sge	10.1	7000	1.5	4.0	4.0	4.0
W Vul	10.1	7000	1.5	4.0	4.0	4.0
W Nor	10.1	7000	1.5	4.0	4.0	4.0
W Sge	10.1	7000	1.5	4.0	4.0	4.0
W Vul	10.1	7000	1.5	4.0	4.0	4.0
W Nor	10.1	7000	1.5	4.0	4.0	4.0
W Sge	10.1	7000	1.5	4.0	4.0	4.0
W Vul	10.1	7000	1.5	4.0	4.0	4.0
W Nor	10.1	7000	1.5	4.0	4.0	4.0
W Sge	10.1	7000	1.5	4.0	4.0	4.0
W Vul	10.1	7000	1.5	4.0	4.0	4.0
W Nor	10.1	7000	1.5	4.0	4.0	4.0
W Sge	10.1	7000	1.5	4.0	4.0	4.0
W Vul	10.1	7000	1.5	4.0	4.0	4.0
W Nor	10.1	7000	1.5	4.0	4.0	4.0
W Sge	10.1	7000	1.5	4.0	4.0	4.0
W Vul	10.1	7000	1.5	4.0	4.0	4.0
W Nor	10.1	7000	1.5	4.0	4.0	4.0
W Sge	10.1	7000	1.5	4.0	4.0	4.0
W Vul	10.1	7000	1.5	4.0	4.0	4.0
W Nor	10.1	7000	1.5	4.0	4.0	4.0
W Sge	10.1	7000	1.5	4.0	4.0	4.0
W Vul	10.1	7000	1.5	4.0	4.0	4.0
W Nor	10.1	7000	1.5	4.0	4.0	4.0
W Sge	10.1	7000	1.5	4.0	4.0	4.0
W Vul	10.1	7000	1.5	4.0	4.0	4.0
W Nor	10.1	7000	1.5	4.0	4.0	4.0
W Sge	10.1	7000	1.5	4.0	4.0	4.0
W Vul	10.1	7000	1.5	4.0	4.0	4.0
W Nor	10.1	7000	1.5	4.0	4.0	4.0
W Sge	10.1	7000	1.5	4.0	4.0	4.0
W Vul	10.1	7000	1.5	4.0	4.0	4.0
W Nor	10.1	7000	1.5	4.0	4.0	4.0
W Sge	10.1	7000	1.5	4.0	4.0	4.0
W Vul	10.1	7000	1.5	4.0	4.0	4.0
W Nor	10.1	7000	1.5	4.0	4.0	4.0
W Sge	10.1	7000	1.5	4.0	4.0	4.0
W Vul	10.1	7000	1.5	4.0	4.0	4.0
W Nor	10.1	7000	1.5	4.0	4.0	4.0
W Sge	10.1	7000	1.5	4.0	4.0	4.0
W Vul	10.1	7000	1.5	4.0	4.0	4.0
W Nor	10.1	7000	1.5	4.0	4.0	4.0
W Sge	10.1	7000	1.5	4.0	4.0	4.0
W Vul	10.1	7000	1.5	4.0	4.0	4.0
W Nor	10.1	7000	1.5	4.0	4.0	4.0
W Sge	10.1	7000	1.5	4.0	4.0	4.0
W Vul	10.1	7000	1.5	4.0	4.0	4.0
W Nor	10.1	7000	1.5	4.0	4.0	4.0
W Sge	10.1	7000	1.5	4.0	4.0	4.0
W Vul	10.1	7000	1.5	4.0	4.0	4.0
W Nor	10.1	7000	1.5	4.0	4.0	4.0
W Sge	10.1	7000	1.5	4.0	4.0	4.0
W Vul	10.1	7000	1.5	4.0	4.0	4.0
W Nor	10.1	7000	1.5	4.0	4.0	4.0
W Sge	10.1	7000	1.5	4.0	4.0	4.0
W Vul	10.1	7000	1.5	4.0	4.0	4.0
W Nor	10.1	7000	1.5	4.0	4.0	4.0
W Sge	10.1	7000	1.5	4.0	4.0	4.0
W Vul	10.1	7000	1.5	4.0	4.0	4.0
W Nor	10.1	7000	1.5	4.0	4.0	4.0
W Sge	10.1	7000	1.5	4.0	4.0	4.0
W Vul	10.1	7000	1.5	4.0	4.0	4.0
W Nor	10.1	7000	1.5	4.0	4.0	4.0
W Sge	10.1	7000	1.5	4.0	4.0	4.0
W Vul	10.1	7000	1.5	4.0	4.0	4.0
W Nor	10.1	7000	1.5	4.0	4.0	4.0
W Sge	10.1	7000	1.5	4.0	4.0	4.0
W Vul	10.1	7000	1.5	4.0	4.0	4.0
W Nor	10.1	7000	1.5	4.0	4.0	4.0
W Sge	10.1	7000	1.5	4.0	4.0	4.0
W Vul	10.1	7000	1.5	4.0	4.0	4.0
W Nor	10.1	7000	1.5	4.0	4.0	4.0
W Sge	10.1	7000	1.5	4.0	4.0	4.0
W Vul	10.1	7000	1.5	4.0	4.0	4.0
W Nor	10.1	7000	1.5	4.0	4.0	4.0
W Sge	10.1	7000	1.5	4.0	4.0	4.0
W Vul	10.1	7000	1.5	4.0	4.0	4.0
W Nor	10.1	7000	1.5	4.0	4.0	4.0
W Sge	10.1	7000	1.5	4.0	4.0	4.0
W Vul	10.1	7000	1.5	4.0	4.0	4.0
W Nor	10.1	7000	1.5	4.0	4.0	4.0
W Sge	10.1	7000	1.5	4.0	4.0	4.0
W Vul	10.1	7000	1.5	4.0	4.0	4.0
W Nor	10.1	7000	1.5	4.0	4.0	4.0
W Sge	10.1	7000	1.5	4.0	4.0	4.0



more massive than about  $0.8 M_{\odot}$  should have long ago died. We here confirm the proposal that this star is the result of a coalescence of two stars in a binary system that had an initial separation smaller than the red giant radius of the more massive star.

Figure 1 gives the work done in each of the 195 zones over each pulsation cycle to cause pulsations in each of the three lowest radial modes. The mass used for this nonadiabatic pulsation analysis is  $1.4 M_{\odot}$ , and the luminosity is the observed 257 suns. The typical population II composition used is the King 1a table with  $Z=0.001$ . Here at 7500K, hotter than NGC 5466 V19, all modes are stable because the radiative damping of the interior (lower zone number) layers is greater than the hydrogen and helium driving in the surface layers.

Figure 2 gives the same plot for the effective temperature of 6500K. In this case it is apparent that the first and second overtone modes are driven more than they are damped, and linear theory predicts growing pulsations. Actually the first overtone is driven the strongest, and at the observed effective temperature of 7000K, probably only the first overtone is unstable to pulsations.

Fig. 1. The work per pulsation cycle to drive pulsations in a NGC 5466 V19 model at 7500K is plotted for each of the 195 zones.





Fig. 2. The work per pulsation cycle to drive pulsations in a NGC 5466 V19 model at 6500K is plotted for each of the 195 zones.



#### REFERENCES

- Becker, S.A., Iban, I., and Tuggle, R.B. (1977). *Ap. J.* **218**, 633.  
 Cox, A.N. (1979). *Ap. J.* **229**, 212.  
 Fernie, J.D. & McGonegal, R. (1983). *Ap. J.* **275**, 732.  
 Kraft, R.P. (1961). *Ap. J.* **134**, 616.  
 Pel, J.W. (1978). *Astr. Ap. Suppl.* **31**, 489.  
 Schmidt, E.G. (1984). *Ap. J.*, in press.  
 Vandenberg, D.A. & Bridges, T.J. (1984). *Ap. J.* **278**, 679.  
 Zinn, R. & King, C.R. (1982). *Ap. J.* **262**, 700.